MEDICAL INFORMATICS IN PERINATOLOGY PROJECT AGUSTINA IN ARGENTINA

DIEGO F. WYSZYNSKI¹, EDUARDO RODAS², PABLO MORICI³, SILVIA DAVEGGIO³, ROBERTO GALLINO FERNANDEZ³, GRACIELA MANETTI⁴, ANGEL TONIETTO³

¹Department of Epidemiology, School of Hygiene and Public Health, The Johns Hopkins University, Baltimore, Maryland, U.S.A. ²Division of Medical Informatics, ³Office of the Director, ⁴Department of Obstetrics and Gynecology, Hospital Zonal General de Agudos Manuel Belgrano, San Martín, Buenos Aires

Summary This paper reviews the development of the field of medical informatics in Latin America. It also describes the preliminary results of a computer-based data management system, named AGUSTINA, which is comprised of maternal and infant data on 6195 deliveries that occurred between June 1990 and December 1995 in a hospital in the surroundings of Buenos Aires, Argentina. These data were fundamental for the instrumentation of preventive community-oriented activities in the area. Finally, this paper describes recommendations for future actions in the area of medical informatics in Latin America.

Keywords: computers, public health, epidemiology, surveillance, statistics

Medical informatics is a field that focuses on the creative use of computers in support of patient care, medical education, and biomedical research¹. Its origins date back to the early 1970s, when the biomedical and computer sciences reached a common path and it was realized that they could complement each other. The development of medical informatics in Latin America, however, is more recent, and irregular across countries. Mexico, Brazil, Argentina, and Cuba took the lead in this region, between 1975 and 1980, by developing the first networks that allowed some of their health care workers access to biomedical information².

The concept of networking is such that individuals "talk" to one another via computers and information is gained and sent via computer technology. Electronic mail (e-mail) is the primary basis for information interchange. Thus, within a

hospital, or among different institutions, one can send notes to individuals electronically by typing a message on a computer and the message is then delivered almost instantaneously to the other person. Messages can be "broadcasted" to several people; e.g., the cardiology department in a hospital or a group of researchers in hospitals all over the world. Bulletin boards can be set up where one logs onto the computer and can find out the events which are occurring within that hospital or group of researchers today. Thus, within a hospital or a field of common interest, the e-mail system markedly improves the speed of information transfer.

With the support of both the World Health Organization (WHO) and the Pan American Health Organization (PAHO), medical personnel of several Latin American countries gathered at national and international meetings, where they exchanged important information and created new networks.

Received: 1-X-1996

Accepted: 24-I-1997

Postal address: Dr. Diego F. Wyszynski, Department of Epidemiology, School of Higiene and Public Health, The Johns Hopkins University, 615 N. Wolfe Street, Baltimore, MD 21205, USA Abbreviations: PCR: perinatal clinical record; CLAP, Centro Latino-Americano de Perinatología; WHO, World Health Organization; PAHO, Pan-American Health Organization; E-Mail, electronic mail; WWW, world wide web; FTP, file transfer protocol

Often, they would go back to their countries to find that, to their despair, no government collaboration could be obtained. The results of some of these efforts are summarized in a document entitled "Health, Information Society and Developing Countries".

Prohibitive telecommunication costs impose a major barrier to the widespread access to on-line services. The communications are largely monopolized by international consortiums in most Latin American countries. Thus, it is not unusual to witness under-utilized software and hardware. Non-profit organizations, state, and academic institutions in the health field struggle to survive in economical systems that are rapidly turning to free-market⁴.

Computer-based information systems (or informatics) and computer-based networking and communications (or telematics) reached the health sector in Argentina in the last ten years. With the collaboration of PAHO, a few groups have been actively seeking, providing and using telematics support to health care research and, to a lesser extent, services. The main purpose of this paper is to present preliminary results of the data obtained using a computer system installed in the Department of Maternity in a hospital located in the surroundings of Buenos Aires, Argentina.

Materials and Methods

The Hospital

The Hospital Zonal General de Agudos «Manuel Belgrano» is located in the city of San Martín, a suburb of the province of Buenos Aires, Argentina. Its programmatic area covers a population of 120,000 inhabitants, embracing large areas of extreme poverty. It has 176 beds, of which 120 are for adults and 56 for children, plus 24 additional maternity beds and 12 others for intensive and intermediate care. This hospital contains 265 medical workers (45% of them nurses) and 81 administrative and maintenance personnel. Additionally, 64 medical residents and 9 post-doctoral fellows are training in this hospital.

Data collection, software and hardware

Beginning in June 1990, the Department of Maternity has collected data using the Perinatal Clinical Record (PCR), developed by the Latin American Perinatology Center (CLAP), in Montevideo, Uruguay, in 1984 ⁵⁻⁸. The PCR records maternal demographic and health information, data on controls and vaccinations during pregnancy, and details on delivery and health status of the newborn. The PCR is administered in a stepwise fashion (Fig. 1) by trained health care workers⁹. The information is then stored on a PC using AGUSTINA (v5.0/1)¹⁰, a data base software designed to store the data obtained using PCR. AGUSTINA includes powerful statistical routines




Fig. 1.- Stepwise administration of Perinatal Clinical Records in the Hospital Manuel Belgrano (modified from9)

and may also be linked to EpiInfo¹¹ for data analysis. Forty-three personal computers were installed in 26 sectors of the hospital, connected to each other by a local area network (LAN).

Results

Between June 1990 and December 1995, there were 6195 registered deliveries (mean: 92 deliveries per month). Ten percent of the women giving birth (*n*=620) were 17 years-of-age or younger and 57 percent (*n*=3531) were out-wedlock, either living with their partners (*n*=2044) or without them (*n*=1487). Mothers who reported no prenatal visits had the highest perinatal mortality (65.5/1,000 liveborns) while mothers with 9 or more prenatal controls had no perinatal deaths. Additionally, mothers who were 35 years-of-age or older had a three-fold increase in perinatal mortality compared to those 18 years-of-age or younger (62.2/1,000 liveborns). Table 1 summarizes perinatal findings obtained using the data base system.

Subrecording was evaluated by cross-checking with other records systems used in the hospital. Additionally, it was possible to determine that maternal height and weight and prenatal visits were the data most likely to be unevenly recorded. Maternal height was missing between 17.82 to 47.44 percent of the times, maternal weight was unrecorded 9.43-34.06 percent of the times, and number of prenatal visits was absent 0.63-33.28 percent of the times, depending on the year.

Discussion

The interaction and the increased performance of computer-based systems have allowed health management officials and the medical community to achieve benefits that may eventually translate into positive actions for the community they serve. By using the AGUSTINA perinatal system, the Department of Maternity has collected the records corresponding to all deliveries performed in the Hospital Manuel Belgrano between June, 1990 and December, 1995, and identified the main needs of its patients. Additionally, the hospital has standardized its recording system and opened the possibility for data exchange with other institutions in Argentina and in Latin America¹².

Data showed that this hospital is covering an extensive programmatic area, comprised mainly of low income and underemployed individuals^{9, 13}. The proportion of young and out-wedlock mothers is remarkably high. Thus, a Pediatric Gynecology Clinic was implemented in collaboration with the Family Planning Clinic, and an intensive outreach program in local schools and health centers was organized and is currently being conducted.

Since the data suggested that mothers who received no prenatal care have the highest perinatal mortality, all hospital primary care physicians were encouraged to identify and educate pregnant women on preventive measures. In addition, perinatal mortality rates showed an increase in aging mothers; therefore, a specialized clinic for

TABLE 1.— Perinatal information gathered using the Perinatal Clinical Records (PCR) and AGUSTINA in Hospital Manuel Belgrano in San Martín, Buenos Aires, Argentina, June 1990 and December 1995 (6,195 registered births)

year	Caesarian incisions (%)	Death during delivery (%)	Low birthweight¹ (%)	Preterm deliveries ² (%)	Late fetal death³ (%)	Early neonatal death4 (%)
1990	23.06	0.21	12.90	15.90	8.5	34.5
1991	23.68	1.53	9.20	11.20	13.5	25.3
1992	26.23	0.19	9.30	10.60	11.8	23.9
1993	28.52	0.18	10.00	12.40	12.8	23.2
1994	26.46	0.00	9.60	10.50	5.2	15.8
1995	22.79	0.00	7.64	9.00	6.7	11.9

'low birthweight: <2500g; ²births before 37 weeks gestation; ³death between 28 weeks of gestation and expected delivery date; ⁴death between birth and the 7th day of life

older mothers was created and implemented. As a consequence of these preventive activities, and supported by other social measures taken by the local authorities, fetal, maternal, and early neonatal death rates have been steadily decreasing in the last years.

In recent years, it has been recognized world wide that poor health of a country affects economies. However, maintaining the health of a community is expensive; the health care expenditure in Argentina represents 8% of the Gross National Product, which is over US \$ 5 billion⁴. The concern produced by these expenditures impose a need for a better use of the existing resources and for the implementation of cost-effective programs in health care. Tools such as computer-based systems, therefore, will be more frequently available in Latin American hospitals and, consequently, previously unrecorded data will become readily accessible for planning and research.

The clinical side of medical informatics spans a broad spectrum, from basic research in how to model medical knowledge to the real-world problems of building a comprehensive computer-based patient record1. In Latin America, the challenge is to develop surveillance report systems to identify the needs and characteristics of the population being served by the various health care providers. AGUSTINA, as presented in this paper, was instituted in a local hospital immersed in a lowclass area of the surroundings of the province of Buenos Aires. The availability of health information in this hospital allows the authorities and professionals to delineate the demographic and epidemiologic characteristics of their patients and to conceive and implement specific measures of disease prevention and control. Additionally, the installation of networking services, such as E-Mail, World Wide Web (WWW), and File Transfer Protocol (FTP), vigorously triggered the enthusiasm for more interinstitutional collaborations.

Just as Homer Warner recommended at the American College of Medical Informatics' (ACMI) Distinguished Lecture, in 1988¹⁴, that medical informatics needs to become an established part of academia, we subscribe to the idea that medical informatics must be a part of all levels of health care. Therefore, hospitals must: 1) establish formal departments of medical informatics where there are now just foci of record-keeping; 2) encourage participation by health care workers from

other departments on the assumption that they will get hooked once they experience the intellectual challenge of this discipline; 3) define training opportunities for health professionals and biomedical students to dedicate time to this field; and 4) establish relationships between industry, government, hospitals, and academic institutions that will lead to support and encouragement of research.

Acknowledgements: We would like to thank Mrs. Denise Wiesch for careful reading of this manuscript. Dr. Wyszynski is supported by an Inter-Personal Agreement (I.P.A.) with the National Institutes of Health, Bethesda, Maryland. We would like to thank all the health care workers in the Hospital Zonal de Agudos Manuel Belgrano who are involved in the program described in this paper.

Resumen

Informática médica en perinatología. Proyecto AGUSTINA en Argentina

Este trabajo revé el desarrollo de la informática médica en América Latina. También describe resultados preliminares de un sistema informático de administración de datos, denominado AGUS-TINA, que incluye información materno-infantil de 6195 partos ocurridos entre junio de 1990 y diciembre de 1995 en un hospital de las afueras de Buenos Aires (Hospital Zonal General de Agudos Manuel Belgrano, San Martín, Pcia. de Buenos Aires). Estos datos fueron fundamentales para la instrumentación de actividades orientadas a la prevención en dicha comunidad. Finalmente, este trabajo propone recomendaciones para acciones futuras en informática médica para el área de América Latina.

References

- Miller PL. Medical informatics in clinical medicine and the biosciences. Nature Med 1995:1:93:94.
- Rienhoff O. Health informatics and telematics in Latin-America: Scenarios and actors. *In:* Health, information society and developing countries, Sosaludicissa M, Levett J, Mandil S, Beales PF (eds.). Amsterdam:IOS Press, 1995.
- Sosa-Iudicissa MC, Levett J, Mandil SH, Beales PF (eds.). Health, information society and developing countries. Amsterdam: IOS Press, 1995.
- Mazzafero VE, Wyszynski DF, Giacomini H. Public Health in Argentina: Facing the Twenty-First Century. J Pub Health Med 1996:18:234-237.

- Schwarcz R, Diaz AG, Fescina R, Rossello JD, Martell M, Tenzer S. Historia clínica simplificada. Publicación Científica del CLAP, Number 1045. Montevideo:CLAP, 1984.
- CLAP. Historia del Sistema Informático Perinatal. Salud Perinatal 1987:2:81-92.
- CLAP. Instructivo para el llenado de la Historia Clínica Perinatal Simplificada Base. Salud Perinatal 1988:S2:1-8.
- Giacomini H, Marconi E. CLAP, Latin American Perinatology Center in Montevideo - Scope and impact. *In:* Health, information society and developing countries, Sosa-Iudicissa M, Levett J, Mandil S, Beales PF (eds.). Amsterdam: IOS Press, 1995.
- Rodas E, Manetti G, Fixman S, López N, San Martín A. Agustina en el Hospital «Manuel Belgrano».

- Medicine 1993:9:79-83.
- San Pedro M. Programa Agustina. Buenos Aires: Fundación Epson, 1990.
- Centers for Disease Control and Prevention. Epilnfo, version 6.03. Atlanta: CDC, 1996.
- CLAP. La Red Perinatal. Salud Perinatal, 1987; 2: 97-99.
- Rodas E, Morici P, Daveggio S, Fernandez-Gallino R, Tonietto A. Computerised systems in the Belgrano Hospital and the promotion of self-governing groups. In: Health, information society and developing countries, Sosa-Iudicissa M, Levett J, Mandil S, Beales PF (eds.). Amsterdam: IOS Press, 1995.
- Warner HR. Medical informatics: a real discipline? JAMA 1995;2:207-214.

Ten of thousands of Americans now regularly telephone "psychics" for information. Do they differ from ancient Romans who sought similar advice from the patterns of animal livers and entrails, or from the voices of oracles?

Decenas de miles de americanos, ahora, telefonean regularmente a "psíquicos" por información. ¿Difieren de los antiguos romanos que buscaban consejo similar de los caracteres del hígado y otras entrañas, o de las voces de los oráculos?

Martin Gardner

When superstition displaces science. Comentario bibliográfico de Behind the Crystal Ball: Magic and Science from Antiquity to the New Age. Anthony Aveni, New York: Random House, 1996. Aparecido en Nature 1997; 385: 405-6.